Draft version February 5, 2008
Preprint typeset using LaTeX style emulateapj v. 10/09/06

arXiv:astro-ph/0607442v3 26 Jul 2007

MERGERS OF BINARY COMPACT OBJECTS

Chang-Hwan Lee, Hong-Jo Park
Department of Physics, Pusan National University, Busan 609-735, Korea
and
Gerald E. Brown
Department of Physics and Astronomy, State University of New York, Stony Brook, NY 11794, USA
Draft version February 5, 2008

ABSTRACT

We work out the effects of hypercritical accretion, which transfers mass from the secondary to the primary (first-born) neutron star (NS) in a binary, showing that the mass of the primary would end up $\gtrsim 0.6 M_\odot$ greater than the secondary NS in contradiction with the very nearly equal masses of the two NS's in binaries measured to date. We discuss that the primary NS evolves into a low-mass black-hole (LMBH) due to the hypercritical accretion. Using a flat distribution in the mass ratio q ($q \equiv M_{\rm secondary}/M_{\rm primary}$), favored by Duquennoy & Mayor (1991), we calculated a ratio of LMBH-NS and NS-NS systems to be ~ 5, in rough agreement with Pinsonneault & Stanek (2006). These authors emphasize the importance of "twins", which we discuss. The two NS's in the twins would be close in mass and further increase the number of mergings.

Subject headings: stars: neutron — binaries: close — stars: evolution — stars: statistics

1. INTRODUCTION

Bethe & Brown (1998) suggested that mergers resulting in short-hard gamma-ray bursts would be mainly those of LMBH-NS binaries, with those of NS-NS binaries down by an order of magnitude from these. The lower number of the latter resulted from the necessity that the two giant progenitors be within 4% of each other in ZAMS mass so that they burned He at the same time. Otherwise the first born pulsar would find itself in the red giant envelope of the companion giant as it evolved and accrete enough matter to go into a black hole (BH). Brown (1995) had already estimated this to be true, based on Chevalier (1993), and proposed the special way that two giants burn He at the same time, in order to avoid the red giant common envelope evolution of the first born pulsar. If the two giants burn He at the same time, the two He envelopes are assumed to go into common envelope evolution, expelling the common envelope matter so that the helium envelope is lost from each star. Braun & Langer (1995) showed that there was not sufficient time for either of the stars to accrete the common envelope He, so that if the two He stars had to be nearly equal in mass, then their progenitor giants must also be. From the Schaller et al. (1992) models for the giant progenitors of the neutron stars we consider, the giants have to be within $\lesssim 4\%$ of each other in mass in order to burn He at the same time. Of the Bethe & Brown (1998) merger rate of 10^{-4} yr^{-1} in our Galaxy, only ~ 0.1, or 10^{-5} yr^{-1} were estimated to be those of binary NS's. A recent detailed calculation by Dewi et al. (2006) gives a merger rate of $0.1 - 12$ Myr^{-1} for Brown's special scenario, the upper end of the calculation in agreement with Bethe & Brown (1998). The latter authors simply estimated that the remaining mergers would be of LMBH-NS binaries since they calculated that when the pulsar went through common envelope with the companion star it accreted $\sim 1 M_\odot$ of matter, enough to send it into a BH. The amount of accretion was corrected downwards $\sim 25\%$ by removal of an approximation of Bethe & Brown (1998) by Belczyński et al. (2002). In the present note we try to make a real calculation of the LMBH-NS binary, NS-NS binary ratio.

Pinsonneault & Stanek (2006) assembled evidence that "Binaries like to be Twins". They showed that a recently published sample of 21 detached eclipsing binaries in the Small Magellanic Cloud can be evolved in terms of a flat mass function containing 55% of the systems and a "twins" population with $q > 0.95$ containing the remainder. All of the binaries had orbital period $P < 5$ days, with primary masses $6.9 M_\odot < M_1 < 27.3 M_\odot$.

The important role of twins is that the two giants are close enough in mass [1] that in Brown (1995) scenario they can evolve into NS-NS binaries, whereas if they are further apart in mass they will evolve into a LMBH-NS binary (Chevalier 1993; Bethe & Brown 1998). Thus the twins may increase the number of NS-NS binaries. We suggest that the resulting number of short hard gamma-ray bursts, which result from the merging of the binaries, which to date are unable to differentiate between the two species, may not be changed much, some of the predicted large excess of LMBH-NS binaries appearing rather as NS-NS binaries. However, because the latter are so much more easy to observe, the role between what we see and what is present will be tightened.

In Sec. 2 we shall show that evolution of binaries with a flat mass evolution does, in agreement with Pinsonneault & Stanek (2006) produce a ratio of (NS+BH)/(NS+NS) systems of 5, for the population that does not contain twins. This is half the order of magnitude ratio found by Bethe & Brown (1998). The lower value results

Electronic address: clee@pusan.ac.kr

---

[1] Pinsonneault & Stanek (2006) used 5% whereas we prefer 4% as will be discussed.



FIG. 1.— Probability density of the formation of NS-NS and NS-LMBH binaries. The x-axis is the ZAMS mass of the progenitors of primary stars in compact binaries. Here we assumed that the accreted pulsars, whose masses are bigger than $1.8M_{\odot}$, went into black holes as in Fig. 2.

from the fact that the secondary in the mass function is not independent of the primary, as we shall see. Taking the non twin binaries to be 55% of the total, the remaining 45% twins, Pinsonneault & Stanek (2006) get a good fit to the detached binaries measured in the Small Magellanic Cloud.

We point out that Belczyński et al. (2002) in their simulation D2 in which the maximum NS mass is $1.5M_{\odot}$ and the mass proximity in the progenitor binaries (to evolve NS's) is taken, like Pinsonneault & Stanek (2006) to be 5%, obtain a ratio of 4 for (BH+NS)/(NS+NS) and would obtain the ratio of 5 had they used our 4% proximity in masses.

In short, there is general agreement amongst the authors quoted above, except that it is not clear how many twins will be left once selection effects are taken into account. Historically large selection effects have been identified (Goldberg et al. 2003; Hogeveen 1992). These will lower the number of twins found by Pinsonneault & Stanek (2006).

In Sec. 3 we summarize the calculations of maximum NS mass obtained in a renormalization group calculation expanding about the chiral restoration fixed point in the vector manifestation of the Harada-Yamawaki hidden local symmetry theory (Brown et al. 2006). The result of $1.5M_{\odot}$ brings us into conflict with the recent measurement of Nice et al. (2005) of $2.1 \pm 0.2M_{\odot}$ for PSR J0751+1807. The lower limit of this NS with 95% confidence level is $1.6M_{\odot}$. We evolve in our model of hypercritical accretion the masses of pulsars and companions in NS-NS binaries, the result of which is that the pulsars would have substantially greater masses than their companions were a $2.1M_{\odot}$ NS star to be stable. We find that our calculated distribution could be made consistent with present observations, the most important of which require pulsar mass to be no greater than $1.8M_{\odot}$. Thus we may have to raise our calculated upper limit $0.2M_{\odot}$ but we believe that the lower end of the mass measurement of J0751+1807 is favored.

2. EVOLUTION OF NON-TWIN BINARIES

In the sample from the Small Magellanic Cloud, Pinsonneault & Stanek (2006) found that the non-twin binaries and the twins are roughly equal (55 % and 45 %, respectively), and the non-twin population can be evolved successfully with a flat mass distribution $dn/dm =$ constant.

Using a flat q ($q \equiv M_2/M_1$)distribution, in which the IMF for the secondary (less massive) star is independent of its mass, $dN/dM_2 = \text{Constant}/M_1$ for a given primary mass M_1, one can obtain the probability of having binaries with initial ZAMS masses within 4%.

$$P_{\rm NS-NS} = \frac{\int_{M_{\rm min}}^{M_{\rm max}} P_{\rm primary}(M_1) \ \left[\int_{\max[M_{\rm min},0.96M_1]}^{M_1} (dN/dM_2)\, dM_2\right] dM_1}{\int_{M_{\rm min}}^{M_{\rm max}} P_{\rm primary}(M_1) \ \left[\int_{M_{\rm min}}^{M_1} (dN/dM_2)\, dM_2\right] dM_1} \tag{1}$$

where $M_{\rm min}$ and $M_{\rm max}$ are the lower and upper limits in ZAMS masses for which neutron star can be formed after supernova explosion, and $P_{\rm primary}$ is the mass function for the primary in binaries. We use the Salpeter mass function as the IMF for the primary,

$$P_{\rm primary}(M_1) \propto M_1^{-2.35}. \tag{2}$$

Note that the lower limit of $0.96M_1$ in the numerator comes from the fact that the mass of the secondary star has to be within 4% in ZAMS mass in order to make the double pulsar.

There is no clear understanding on the relation between the pulsar masses and their progenitors. We gave arguments that the Hulse-Taylor pulsar 1913+16 was evolved from a $20M_{\odot}$ ZAMS giant (Bethe et al. 2007). (See

Burrows & Woosley 1986.) The companion in J1756−2251 of $1.18^{+0.03}_{-0.02}$ is the lowest measured mass in NS binaries, and we take it to have originated from a $10M_\odot$ giant. Our interval in which binary NS's can be formed is taken to be $10 - 20M_\odot$ as in Bethe & Brown (1998).

From eq. 2, with given assumptions, we finds that NS-NS binaries should be formed 16% of the time, resulting in 5.3 times more NS-LMBH binaries, which are harder to be observed, than NS-NS binaries. The estimated probability distribution is summarized in Fig. 1. From the figure, one can see that more NS-NS binaries can be formed with lower ZAMS mass primary stars, while LMBH-NS binaries mainly come from binaries with higher ZAMS mass primary stars (mainly because the possibility of forming NS stars out of secondary stars are higher).

By requiring stars to be within 5% of each other in mass to burn He at the same time, Belczyński et al. (2002) get a ratio of 4 to 1 for LMBH-NS binaries over NS-NS binaries. From eq. (2) with 5%, we obtained 20% chance of making NS-NS binaries, giving 4 times more LMBH-NS binaries than NS-NS binaries, which are consistent with Belczyński et al. (2002) within 10% uncertainty. So not only does the flat distribution fit the non-twin binaries in the Small Magellanic Cloud but it reproduces Belczyński's result for the same assumed proximity of 5% in mass. (Note that there can be extra uncertainties in this direct comparison because different evolutionary phases in different approaches can alter the final ratio.)

The hypercritical accretion plays a central role in the result of Belczyński et al. (2002) as well as ours and the agreement between our results and their is perhaps better than would be expected in such a complex situation. We know hypercritical accretion to be present, because the trapping of photons and subsequent adiabatic inflow is based on the same physics, but with quite different parameters, as the trapping of neutrinos in the infall of matter in the collapse of large stars. A rough estimate of the correction to the original Bethe & Brown (1995) result, which neglected the compact object mass compared with the varying giant mass, gives the $\sim 25\%$ decrease to get from the Bethe & Brown (1995) results to those of Belczyński et al. (2002).

The supernova explosion in a binary of two He stars to produce a double neutron star were carried out by Bethe & Brown (1998) in a simple schematic way with a simple physical description of the survival rates, i.e., of the binary staying together after each explosion. Computer calculations by Wettig & Brown (1996) showed the Bethe & Brown (1998) assumption to be accurate to $\leq$ 10%, slightly greater than with the formula of Portegies Zwart & Yungelson (1998). In principle the merging of neutron stars and black holes could be very different because initially the mass transfer from neutron star into black hole is stable, so the black hole can "bounce" away from the neutron star (Portegies Zwart & Yungelson 1998). The regularities in this are usually assumed to be broken up by the gravitational interactions, etc. and the binaries are assumed to merge by some kind of Roche Lobe overflow. Thus, although the manner of merging may be quite different for binary neutron stars and a neutron star and low-mass black hole, we believe that the ratio of mergings will be given by the population synthesis.

3. BINARY NEUTRON STARS VS LMBH-NS BINARIES

Kaon condensation has been suggested as a possible mechanism which can reduce the maximum mass of neutron stars (Bethe & Brown 1995). Recently, a theoretical calculation of strangeness condensation has been carried out by expanding about the fixed point of the Harada-Yamawaki Vector Manifestation (Brown et al. 2006). In the strangeness condensation, which takes place at a density not much less than that of chiral restoration, the fixed point, the in-medium K^- mass $m^\star_{K^-}$ has been brought down to the value of the electron chemical potential μ_e, so that the highly degenerate fermionic electrons collapse into a zero-momentum Bose condensate of K^- mesons. Only mesons and constituent quarks are left at densities close to the chiral restoration density $n_{\chi SR}$. Expanding about the fixed point eliminates the uncertainties introduced by many investigators about the role of strange mesons and the role of the explicit symmetry breaking in the strangeness sector. In fact, Brown et al. (2006) show that the K^- meson behaves like the nonstrange mesons, ρ, π, σ, A_1 in that its mass goes to zero as $n \to n_{\chi SR}$. However, the criterion for kaon condensation is much more easily fulfilled, because the *in-medium* mass $m^\star_{K^-}$ has only to come down a little more than half way to zero; i.e., to the electron chemical potential μ_e, for the phase transition to take place.

In Brown et al. (2006) interactions between the bosons in the condensate were ignored. They would not be expected to be large because the K^- mesons at condensation have a density of only ~ 0.05 fm $^{-3}$, an order of magnitude less than the neutron density at kaon condensation. Thus, the K^- in the condensation are very diffuse. We can make this more quantitative by calculating their interaction. The optical model potential from the other K^--mesons by a given K^- is

$$V_{\rm opt} = \frac{4\pi a}{(M_{K^-})_{\rm reduced}} n_{K^-} \tag{3}$$

where a is the K^--K^- scattering length and the reduced mass is used. The a can be obtained[2] from the $I = 2$ π-π scattering length by changing M_π to M_{K^-}, giving

$$a = -\frac{M_{K^-}}{16\pi f_\pi^2}. \tag{4}$$

Thus

$$V_{\rm opt} = n_{K^-}/2f_\pi^2 \tag{5}$$

---

[2] We are grateful to Christof Hanhart for this information.



FIG. 2.— Masses of primary compact stars with and without accretion during H red giant stage of secondary star (refer Table 1 for the details). Note that the 84% corresponds to $M_{\rm Compact\ Star,primary} > 1.8 M_\odot$. There is uncertainty in the final primary compact star masses due to the extra mass accretion, $\sim 0.1 M_\odot - 0.2 M_\odot$, during He giant stage. This may increase the primary compact star masses for both 'with-' and 'without-accretion'. Note that with our maximum neutron star mass of $1.8 M_\odot$, all primary compact stars with accretion would go into low-mass black holes.

where f_π is the pion decay constant $f_\pi \sim 90$ MeV or one can put in higher-order effects and use $f_K \simeq 108$ MeV. The latter, together with

$$n_{K^-} = 0.048 \text{fm}^{-3} \tag{6}$$

at kaon condensation would give

$$V_{\rm opt} = 16 \text{MeV}, \tag{7}$$

sufficiently negligible compared with the ~ 250 MeV in going from M_{K^-} to $M^\star_{K^-} \sim M_{K^-}/2$ at kaon condensation. Thus, we stick by our calculation in Brown et al. (2006). None the less, in the light of the measurement of the mass of PSR 0751+1802 (Nice et al. 2005), it is very interesting to see how far the maximum neutron star mass can be pushed up without the pattern of masses in neutron star binaries running into conflict with observation. Later we shall see that there is strong evidence of a NS more massive than $1.5 M_\odot$ existing, although we discuss that the limit can be extended to $1.8 M_\odot$ without destroying the pattern of accurately measured NS's to date.

In the original Bethe & Brown (1998) calculation of hypercritical accretion the common envelope evolution was carried out analytically, using Newton's and Kepler's laws and the coefficient of dynamical friction $c_d = 6$. The equations could be solved analytically only if the NS mass M_{NS}, the smallest of the masses, was set equal to zero. This approximation was removed by Belczyński et al. (2002). Incorporating their corrections we find that an originally $1.4 M_\odot$ pulsar will have accreted $0.75 M_\odot$ in the H red giant stage. This is enough to send it into a black hole, and this is our correction of the conventional model which ignores the accretion.

Lower mass He stars have to burn hotter than the higher mass ones because of the loss of energy through the surface. Ordinarily the range of $M_{He} = 2 - 4 M_\odot$ is chosen as a limit below which they begin a red giant phase. We believe from the equality of masses in B1534+12 that not much He can have been transferred in the pulsar, He-star stage of the evolution. We, therefore, introduce He red giant evolution below the ZAMS mass of $\sim 15 M_\odot$ for this binary. The double pulsar with companion mass $1.25 M_\odot$ certainly has been recycled by mass deposited by its companion during its He red giant stage. Whereas we calculate the hypercritical accretion in He red giant to be $0.2 M_\odot$ (Bethe et al. 2007) the actual difference in J0737−3039 between pulsar and companion mass is $0.09 M_\odot$. However, the $0.2 M_\odot$ is roughly correct for J1756−2251. The He wind is quite uncertain. In fact, Belczyński & Kalogera (2001) derived a way in which no mass was accreted, similar to the Brown (1995) double He star scenario, which expelled the H envelopes, except that now the He envelopes from the two stars meet and are expelled. For other possible evolutionary scenarios of NS-NS binaries, please refer Yamaoka et al. (1993); Fryer & Kalogera (1997); Arzoumanian et al. (1999); Francischelli et al. (2002); Dewi & Pols (2003); Willems et al. (2004); Thorsett et al. (2005).

Using the four accurately measured companion masses in relativistic binary NS's [3] we construct the final pulsar masses as in Fig. 2 and Table 1. Our strategy is to calculate the mass accreted on the primary pulsar in the H red giant stage of secondary star. There can be an extra accretion in the He red giant stage, which will increase the final

---

[3] We could easily also fit in 2127+11C, but it is generally thought to have been made by recombination of NS's formed in different binaries.

TABLE 1

Calculated accretion onto the pulsar during H red giant stage of secondary star. Masses are given in $M_\odot$. Note that for a given secondary star mass $M_{\rm ZAMS,secondary}$, the primary star can have masses in the range $M_{\rm ZAMS,secondary} < M_{\rm ZAMS,primary} < M_{\rm max} = 20 M_\odot$. However, in order to have mass accretion during H red giant stage, the mass difference has to be bigger than 4%. So, in the second column, for the estimation of the lower limit of the accretion, we take $M^{\rm min}_{\rm ZAMS,primary} = M_{\rm ZAMS,secondary}/0.96$. In the third column, we take $M^{\rm max}_{\rm ZAMS,primary} = 20 M_\odot$ for the estimation of the upper limit. $M_{\rm C,i}$ is the initial primary compact star mass and $M_{\rm C,f}$ is the final mass following accretion. The He core mass of giant star is assumed to be $M_{\rm He} = 0.08 (M_{\rm Giant}/M_\odot)^{1.45} M_\odot$. Note that there can be extra mass accretion during He giant stage, $\Delta M_{\rm C,f} = 0.1 M_\odot - 0.2 M_\odot$, therefore, $M_{\rm C,f}$ in this table can be treated as a lower limit.

Secondary star		Primary star; Minimum mass			Primary star; Maximum mass		
$M_{\rm ZAMS}$	$M_{\rm NS}$	$M^{\rm min}_{\rm ZAMS}$	$M_{\rm C,i}$	$M_{\rm C,f}$	$M^{\rm max}_{\rm ZAMS}$	$M_{\rm C,i}$	$M_{\rm C,f}$
20	1.40	—	—	—	—	—	—
19	1.39 (B1913+16)	19.8	1.40	2.25	20	1.40	2.25
18	1.38	18.8	1.39	2.21	20	1.40	2.23
17	1.36	17.7	1.37	2.17	20	1.40	2.20
16	1.35	16.7	1.36	2.13	20	1.40	2.18
15	1.34 (B1534+12)	15.6	1.35	2.08	20	1.40	2.15
14	1.32	14.6	1.33	2.03	20	1.40	2.12
13	1.30	13.5	1.31	1.98	20	1.40	2.09
12	1.28	12.5	1.29	1.93	20	1.40	2.06
11	1.25 (J0737−3039B)	11.5	1.26	1.86	20	1.40	2.02
10	1.18 (J1756−2251)	10.4	1.22	1.78	20	1.40	1.99

TABLE 2

Compilation of NS-NS binaries (Lattimer & Prakash 2007). Individual references are (a) Thorsett & Chakrabarty 1999, (b) Lyne et al. 2004, and (c) Faulkner et al. 2004.

Object	Mass ($M_\odot$)	Companion Mass ($M_\odot$)	References
J1518+49	$1.56^{+0.13}_{-0.44}$	$1.05^{+0.45}_{-0.11}$	(a)
B1534+12	$1.3332^{+0.0010}_{-0.0010}$	$1.3452^{+0.0010}_{-0.0010}$	(a)
B1913+16	$1.4408^{+0.0003}_{-0.0003}$	$1.3873^{+0.0003}_{-0.0003}$	(a)
B2127+11C	$1.349^{+0.040}_{-0.040}$	$1.363^{+0.040}_{-0.040}$	(a)
J0737−3039A	$1.337^{+0.005}_{-0.005}$	$1.250^{+0.005}_{-0.005}$ (J0737−3039B)	(b)
J1756−2251	$1.40^{+0.02}_{-0.03}$	$1.18^{+0.03}_{-0.02}$	(c)

primary compact star mass. We also neglected mass transfer by He winds in the pulsar, He star binary which precedes the NS-NS stage, but this is likely to be small because of the propeller effect (Francischelli et al. 2002). For the relation between the pulsar and progenitor masses, we assumed

$$M_{\rm Pulsar} = (1.40 M_\odot - 1.18 M_\odot) \times \left[\frac{M_{\rm ZAMS} - 10 M_\odot}{20 M_\odot - 10 M_\odot} \right]^{0.5} + 1.18 M_\odot. \tag{8}$$

Note that this approximation is based on the known pulsar masses, as in Table 2, and the pulsar and progenitor mass relations, which we discussed above and marked in Table 1.

Using the Bethe & Brown (1998) hypercritical accretion corrected by Belczyński et al. (2002) we arrive at Fig. 2 and Table 1, giving the estimated final primary compact star masses. Note that there can be extra mass accretion during He giant stage, $0.1 M_\odot - 0.2 M_\odot$, therefore, final compact star masses in Fig. 2 and Table 1 can be treated as a lower limit.

In Table 2 we list the known relativistic NS-NS binaries. The masses of the two least massive binaries J0737−3039A, B and J1756−2251 are slightly different because of mass transfer during the He red giant stage, but certainly the near equality of masses within a binary is completely different from that in Table 1. Thus, we conclude that the pulsar must have evolved into a BH in most of the cases, leaving only those from double He star burning; i.e., those in which the two NS's are within 4% of each other in mass.

We note that the masses of NS's evolved in binaries with white dwarfs by Tauris & Savonije (1999) are similar to those in the column labeled M_f in our Table 1, 2/3 of their final masses being greater than $2 M_\odot$. Indeed, their NS mass for J0751+1807 is $2.16 M_\odot$, to compare with the measured mass of $2.1 \pm 0.2 M_\odot$ (Nice et al. 2005) and companion mass is $0.181 M_\odot$ to compare with the Nice et al. (2005) $0.191 \pm 0.015 M_\odot$. The Nice et al. (2005) values here are with 68% confidence, whereas they are $2.1^{+0.4}_{-0.5} M_\odot$ for the NS and $0.191^{+0.033}_{-0.029}$ for the white dwarf with 95% confidence.

Bethe & Brown (1995) obtained a maximum mass for the compact core of 1987A of $1.56 M_\odot$, assuming that it had gone into a black hole, from the $\sim 0.075 M_\odot$ of Ni production. From our result in Fig. 2, the upper limit on the maximum NS mass can be pushed up to $\sim 1.8 M_\odot$. Our maximum NS mass, $\lesssim 1.8 M_\odot$, cannot be reconciled with the observations. However, it is noteworthy that only one of the dozen or so NS, white-dwarf binaries observed exceeds

TABLE 3
Observability premiums of unrecycled and recycled pulsars.

Unrecycled Pulsar	Π
B2303+46	1.26
J1141−6545	0.77
Recycled Pulsar	
J2145−0750	1667
J1022+1001	1190
J1157−5112	159
J1453−5846	164
J1435−6100	2127
J1756−5322	?

our limit, whereas most of Tauris & Savonije (1999) do. The Nice et al. (2005) error will be improved with further observation. It will be interesting to see how their results evolve.

4. COMMON ENVELOPE EVOLUTION IN NS-COWD BINARIES

Our scenario that in NS-NS binary evolution the pulsar goes into a BH if common envelope evolution takes place while the companion is red giant has been challenged in the literature, because it is known that a number of NS, carbon-oxygen white dwarfs (COWD) PSR 1157−5112, J1757−5322, J1435−6100, J1454−5846, J1022+100, and J2145−750 have survived common envelope evolution, from the fact that the NS is strongly recycled. At least two unrecycled pulsars, B2303+46 and J1141−6545 have been observed.

The behavior of the pulsar magnetic field is crucial for the observability times of NS's. Van den Heuvel (1994b) has pointed out that NS's formed in strong magnetic fields $10^{12} - 5\times10^{12}$ Gauss, spin down in a time $\tau_{\rm sd} \sim 5\times10^6$ years and then disappear into the graveyard of NS's. The pulsation mechanism requires a minimum voltage from the polar cap, which can be obtained from $B_{12}/P^2 \gtrsim 0.2$ with $B_{12} = B/10^{12}$ G and P in seconds. As a result, the relative time that the pulsar is observable is proportional to the observability premium Π (Wettig & Brown 1996), the ratio of observability for a pulsar with magnetic field B to one of $B = 10^{12}$ G,

$$\Pi = 10^{12}{\rm G}/B \tag{9}$$

where B is the magnetic field strength of the pulsar, and the spin down depends only on B.

Since the magnetic fields B of recycled pulsars have been brought down substantially by accretion, presumably from their companions, as found phenomenologically by Taam & van den Heuvel (1986), the pulsar that have had a lot of recycling, and, therefore lower B, can be observed for a much longer time than "fresh" pulsars that have not been recycled.

We plot the observability premium of the unrecycled and recycled pulsars separately in Table 3. We see from Table 3 that the chance of observing a recycled binary, because of the longer lifetime of the pulsar, is at least two orders of magnitude greater than observing an unrecycled pulsar. We interpret this as meaning that almost all of the recycled pulsars go into LMBH-NS binaries in common envelope evolution, but there are a few exceptions.

The discussion by Van den Heuvel (1994a) of 2145−0750 is illuminating as to how the NS binary might survive. He requires an efficiency for expelling the envelope of a factor of several greater than usual in common envelope binary evolution (Webbink 1984), interpreting this as meaning that agencies other than the dynamical friction are helping to expel the envelope. The use of efficiencies greater than unity is often employed in order to include phenomenologically additional effects that help the binary survive common envelope evolution (Iben & Livio 1993; De Kool 1990; Yungelson et al. 1993). We know that in the asymptotic giant branch the envelope is lost, in any event, as the originally main sequence star evolves. Possibly the strong gravitational field of the NS helps in expelling the envelope. In any case it is clear that the most likely fate of a NS-COWD which has gone through common envelope evolution is that the NS accretes enough matter to go in a LMBH.

Our assumption is that survival probabilities for the double neutron star formation and NS-LMBH formation are the same. Unfortunately, this cannot be checked because it has not been possible to distinguish in the short-hard gamma-ray bursts between the two types of events (Nakar 2007).

5. CONCLUSION

The work of Pinsonneault & Stanek (2006) on the 21 detached binaries in the Small Magellanic Cloud suggests that there are about as many giant stars in twins as in the non-twin population, although the number in twins will be decreased by selection effects which were not taken into account. The twins are within 5% of each other in ZAMS mass and will therefore burn He at the same time so that binaries with sufficiently massive stars ($\gtrsim$ ZAMS $10M_\odot$) will evolve into NS-NS binaries.

These will add, in the merging of compact binaries to the Bethe & Brown (1998) LMBH-NS binary mergings. Neglecting the contribution from twins, these are shown to increase the NS-NS binary mergers by a factor of 6, down by a factor of 2 from the rough estimate of Bethe & Brown (1998), when evolved with a flat mass function for the companion star. The latter is shown to favor binaries of low mass NS, like the NS-NS binary, because the companion mass is correlated with the pulsar mass by its necessity of being less massive.

We reconsidered the maximum mass determination of $1.5M_\odot$ for a NS, first proposed by Bethe & Brown (1995) and Brown & Bethe (1994), in the light of the measurement of the $2.1 \pm 0.2M_\odot$ mass in PSR 0751+1807. We show that our maximum NS mass could be raised to $1.8M_\odot$, within 95% confidence value of Nice et al. (2005), but that increasing it more would lead to the pulsars of higher mass than $1.8M_\odot$ remaining stable following hypercritical accretion during the red giant evolution of the companion (which is necessarily less massive than the pulsar at formation times of the two NS's). Thus the pattern of NS-NS binaries would chiefly consist of pulsar which are $\sim 50\%$ massive than companion NS, which is not seen.

ACKNOWLEDGMENTS

C.H.L. was supported by the Korea Research Foundation Grant funded by the Korean Government(MOEHRD, Basic Research Promotion Fund) (KRF-2005-070-C00034). G.E.B. was supported in part by the US Department of Energy under Grant No. DE-FG02-88ER40388.

REFERENCES

Arzoumanian, Z., Cordes, J.M., and Wasserman, I. 1999, ApJ, 520, 696.
Belczyński, K. and Kalogera, V. 2001, ApJ, 550, L183.
Belczyński, K., Kalogera, V., and Bulik, T. 2002, ApJ, 572, 407.
Bethe, H.A. and Brown, G.E. 1995, ApJ, 445, L129.
Bethe, H.A. and Brown, G.E. 1998, ApJ, 506, 780.
Bethe, H.A., Brown, G.E., and Lee, C.-H. 2007, Physics Reports, 442, 5.
Braun, H. and Langer, N. 1995, A&A, 297, 483.
Brown, G.E. 1995, ApJ, 440, 270.
Brown, G.E. and Bethe, H.A. 1994, ApJ, 423, 659.
Brown, G.E., Lee, C.-H., Park, H.-J., and Rho, M. 2006, Phys. Rev. Lett. 96, 062303.
Burrows, A. and Woosley, S.E. 1986, ApJ, 308, 680.
Chevalier, R.A. 1993, ApJ, 411, L33.
De Kool, M. 1990, ApJ, 358, 189.
Dewi, J.D.M., Podsiadlowski, Ph., and Sena, A. 2006, MNRAS, 368, 1742.
Dewi, J.D.M. and Pols, O.R. 2003, MNRAS, 344, 629.
Duquennoy, A. and Mayor, M. 1991, A&A, 248, 485.
Faulkner et al. 2004, ApJL, 618, L119.
Francischelli, G., Wijers, R.A.M.J. and Brown, G.E. 2002, ApJ, 565, 471.
Fryer, C. and Kalogera, V. 1997, ApJ, 489, 244.
Goldberg, D., Mazeh, T., and Latham, D. 2003, ApJ, 591, 397.
Harada, M. and Yamawaki, K. 2003, Phys. Rep. 381, 1.
Hogeveen, S.J. 1992, Astrophys and Space Science, 194, 143; ibid 195, 359, ibid 196, 299.
Iben, I. and Livio, M. 1993, PASP, 105, 1373.
Lattimer, J. and Prakash, M. 2007, Physics Reports, 442, 166.
Lyne, A.G., et al. 2004, Science, 303, 1153.
Nakar, E. 2007, Physics Reports, 442, 166.
Nice, D.J., et al. 2005, ApJ, 634, 1242.
Pinsonneault, M.H. and Stanek, K.Z. 2006, ApJ, 639, L67.
Portegies Zwart, S.F. and Yungelson, L.R. (1998), A&A, 332, 173.
Schaller, G., Schaerer, D., Meynet, G., Maeder, A. 1992, A&A, Suppl. Series, 96, 269.
Taam, R.E. and van den Heuvel, E.P.J. 1986, ApJ, 305, 235.
Tauris, T.M. and Savonije, G.J. 1999, A&A, 350, 928.
Thorsett, S.E. and Chakrabarty, D. 1999, ApJ, 512, 288.
Thorsett, S.E., Dewey, R.J., and Stairs, I.H. 2005, ApJ, 619, 1036.
Yungelson, L.R., Tutukov, A.V. and Livio, M. 1993, ApJ, 418, 794.
Van den Heuvel, E.P.J. 1994a A&A, 291, L39.
Van den Heuvel, E.P.J. 1994b, "Interacting Binaries", Lecture notes of the 22nd Advanced Course of the Swiss Society for Astronomy and Astrophysics (SSAA), held at Saas-Fee, Switzerland, 1992, Berlin-Springer, edited by H. Nussbaumer and A. Orr.
Webbink, R.G. 1984, ApJ, 277, 355.
Wettig, T. and Brown, G.E. 1996, New Astronomy, 1, 17.
Willems, B., Kalogera, V., and Henninger, M. 2004, ApJ, 616, 414.
Yamaoka, H., Shigeyama, T., and Nomoto, K. 1993, A&A, 267, 433.